

Product datasheet for **RC202533**

CXCL14 (NM_004887) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: CXCL14 (NM_004887) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: CXCL14
Synonyms: BMAC; BRAK; KEC; KS1; MIP-2g; MIP2G; NJAC; SCYB14
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC202533 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGTCCCTGCTCCACGCCGCGCCCTCCGGTCAGCATGAGGCTCCTGGCGGCCGCTGCTCCTGCTGC
TGCTGGCGCTGTACACCGCGGTGTGGACGGTCCAAATGCAAGTGTCCCGGAAGGGACCAAGATCCG
CTACAGCGACGTGAAGAAGCTGGAATGAAGCCAAAGTACCCGACTGCGAGGAGAAGATGGTTATCATC
ACCACCAAGAGCGTGTCCAGGTACCGAGGTACGAGACTGCCTGCACCCCAAGCTGCAGAGCACCAAGC
GCTTCATCAAGTGTACAACGCCTGGAACGAGAAGCGCAGGGTCTACGAAGAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC202533 protein sequence
Red=Cloning site Green=Tags(s)

MSLLPRRAPPVSMRLLAALLLLLLLALYARVDGSKCKCSRKGPKIRYSVKKLEMPKYPHCEEKMVII
TTKSVSRYRGQEHCLHPKLQSTKRFIKWYNWNEKRRVYEE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6143_e12.zip

Restriction Sites: SgfI-MluI



[View online »](#)

Cloning Scheme:


ACCN: NM_004887

ORF Size: 333 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

| | |
|-------------------------------|---|
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | NM_004887.4 |
| RefSeq Size: | 1989 bp |
| RefSeq ORF: | 300 bp |
| Locus ID: | 9547 |
| UniProt ID: | O95715 |
| Cytogenetics: | 5q31.1 |
| Protein Families: | Druggable Genome, Secreted Protein, Transmembrane |
| Protein Pathways: | Chemokine signaling pathway, Cytokine-cytokine receptor interaction |
| MW: | 13.1 kDa |
| Gene Summary: | This antimicrobial gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation. [provided by RefSeq, Sep 2014] |

Product images:



Circular map for RC202533



Western blot validation of overexpression lysate (Cat# [LY401525]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC202533 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).