

## Product datasheet for RC202557L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Nck beta (NCK2) (NM\_003581) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Nck beta (NCK2) (NM\_003581) Human Tagged ORF Clone Lentiviral Particle

Symbol: Nck beta

Synonyms: GRB4; NCKbeta

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_003581 **ORF Size:** 1140 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202557).

OTI Disclaimer:

Sequence:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 003581.2</u>

 RefSeq Size:
 2517 bp

 RefSeq ORF:
 1143 bp

 Locus ID:
 8440

 UniProt ID:
 043639

 Cytogenetics:
 2q12.2

**Protein Families:** Druggable Genome

SH2, SH3





## Nck beta (NCK2) (NM\_003581) Human Tagged ORF Clone Lentiviral Particle - RC202557L4V

**Protein Pathways:** Axon guidance, ErbB signaling pathway, Pathogenic Escherichia coli infection, T cell receptor

signaling pathway

**MW:** 42.9 kDa

**Gene Summary:** This gene encodes a member of the NCK family of adaptor proteins. The protein contains

three SH3 domains and one SH2 domain. The protein has no known catalytic function but has been shown to bind and recruit various proteins involved in the regulation of receptor protein tyrosine kinases. It is through these regulatory activities that this protein is believed

to be involved in cytoskeletal reorganization. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]