

## Product datasheet for RC202650L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RING1 (NM\_002931) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: RING1 (NM\_002931) Human Tagged ORF Clone Lentiviral Particle

Symbol: RING1

Synonyms: RING1A; RNF1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_002931

 ORF Size:
 1218 bp

**ORF Nucleotide** 

1210 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC202650).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002931.3

RefSeq Size: 1744 bp
RefSeq ORF: 1221 bp
Locus ID: 6015
UniProt ID: Q06587
Cytogenetics: 6p21.32
Domains: RING

**Protein Families:** Druggable Genome, Transcription Factors





ORIGENE

**MW:** 42.4 kDa

**Gene Summary:** This gene belongs to the RING finger family, members of which encode proteins

characterized by a RING domain, a zinc-binding motif related to the zinc finger domain. The gene product can bind DNA and can act as a transcriptional repressor. It is associated with the multimeric polycomb group protein complex. The gene product interacts with the polycomb group proteins BMI1, EDR1, and CBX4, and colocalizes with these proteins in large nuclear domains. It interacts with the CBX4 protein via its glycine-rich C-terminal domain. The gene maps to the HLA class II region, where it is contiguous with the RING finger genes FABGL

and HKE4. [provided by RefSeq, Jul 2008]