

Product datasheet for RC202669L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nucleoside Diphosphate Kinase 7 (NME7) (NM_197972) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nucleoside Diphosphate Kinase 7 (NME7) (NM_197972) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Nucleoside Diphosphate Kinase 7

Synonyms: CFAP67; MN23H7; NDK 7; NDK7; nm23-H7

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_197972 **ORF Size:** 1131 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC202669).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 197972.1</u>

 RefSeq Size:
 1625 bp

 RefSeq ORF:
 1023 bp

 Locus ID:
 29922

 UniProt ID:
 Q9Y5B8

 Cytogenetics:
 1q24.2

Protein Families: Druggable Genome





Nucleoside Diphosphate Kinase 7 (NME7) (NM_197972) Human Tagged ORF Clone Lentiviral Particle - RC202669L4V

Protein Pathways: Metabolic pathways, Purine metabolism, Pyrimidine metabolism

MW: 42.5 kDa

Gene Summary: This gene encodes a member of the non-metastatic expressed family of nucleoside

diphosphate kinases. Members of this family are enzymes that catalyzes phosphate transfer from nucleoside triphosphates to nucleoside diphosphates. This protein contains two kinase domains, one of which is involved in autophosphorylation and the other may be inactive. This protein localizes to the centrosome and functions as a component of the gamma-tubulin ring complex which plays a role in microtubule organization. Mutations in this gene may be associated with venous thromboembolism. Alternate splicing results in multiple transcript

variants. [provided by RefSeq, Sep 2016]