

Product datasheet for RC202703L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caveolin 2 (CAV2) (NM_001233) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Caveolin 2 (CAV2) (NM_001233) Human Tagged ORF Clone Lentiviral Particle

Symbol: Caveolin 2

Synonyms: CAV

Mammalian Cell None

Selection:

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_001233 **ORF Size:** 486 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC202703).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001233.3

 RefSeq Size:
 3332 bp

 RefSeq ORF:
 489 bp

 Locus ID:
 858

 UniProt ID:
 P51636

 Cytogenetics:
 7q31.2

Domains: Caveolin

Protein Families: Druggable Genome, Transmembrane





Protein Pathways: Focal adhesion

MW: 18.1 kDa

Gene Summary: The protein encoded by this gene is a major component of the inner surface of caveolae,

small invaginations of the plasma membrane, and is involved in essential cellular functions, including signal transduction, lipid metabolism, cellular growth control and apoptosis. This protein may function as a tumor suppressor. This gene and related family member (CAV1) are located next to each other on chromosome 7, and express colocalizing proteins that form a stable hetero-oligomeric complex. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. Additional isoforms resulting from the use of alternate in-frame translation initiation codons have also been described, and shown to have

preferential localization in the cell (PMID:11238462). [provided by RefSeq, May 2011]