

Product datasheet for RC202715L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NDUFA2 (NM_002488) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: NDUFA2 (NM_002488) Human Tagged ORF Clone Lentiviral Particle

Symbol: NDUFA2

Synonyms: B8; CD14; CIB8; MC1DN13

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 002488

ORF Size: 297 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202715).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002488.2</u>

 RefSeq Size:
 726 bp

 RefSeq ORF:
 300 bp

 Locus ID:
 4695

 UniProt ID:
 043678

 Cytogenetics:
 5q31.3

Domains: L51_S25_CI-B8





NDUFA2 (NM_002488) Human Tagged ORF Clone Lentiviral Particle - RC202715L3V

Protein Pathways: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation,

Parkinson's disease

MW: 10.9 kDa

Gene Summary: The encoded protein is a subunit of the hydrophobic protein fraction of the

NADH:ubiquinone oxidoreductase (complex 1), the first enzyme complex in the electron transport chain located in the inner mitochondrial membrane, and may be involved in regulating complex I activity or its assembly via assistance in redox processes. Mutations in this gene are associated with Leigh syndrome, an early-onset progressive neurodegenerative disorder. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May

2010]