

Product datasheet for RC202716L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL18 (NM_001562) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL18 (NM 001562) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL18

Synonyms: IGIF; IL-1g; IL-18; IL1F4

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 001562

ORF Size: 579 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202716).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001562.2

 RefSeq Size:
 1163 bp

 RefSeq ORF:
 582 bp

 Locus ID:
 3606

 UniProt ID:
 Q14116

 Cytogenetics:
 11q23.1

Protein Families: Druggable Genome, Secreted Protein



IL18 (NM_001562) Human Tagged ORF Clone Lentiviral Particle - RC202716L1V

Protein Pathways: Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, NOD-like receptor

signaling pathway

MW: 22.3 kDa

Gene Summary: The protein encoded by this gene is a proinflammatory cytokine of the IL-1 family that is

constitutively found as a precursor within the cytoplasm of a variety of cells including macrophages and keratinocytes. The inactive IL-18 precursor is processed to its active form by caspase-1, and is capable of stimulating interferon gamma production, and of regulating both T helper (Th) 1 and Th2 responses. This cytokine has been implicated in the injury of different organs, and in potentially fatal conditions characterized by a cytokine storm. In humans, IL-18 gene is located on chromosome 11. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Aug 2020]