

Product datasheet for RC202736L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDK7 (NM_001799) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CDK7 (NM_001799) Human Tagged ORF Clone Lentiviral Particle

Symbol: CDK7

Synonyms: CAK; CAK1; CDKN7; HCAK; MO15; p39MO15; STK1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_001799 **ORF Size:** 1038 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202736).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001799.2

RefSeq Size: 1534 bp
RefSeq ORF: 1041 bp
Locus ID: 1022
UniProt ID: P50613
Cytogenetics: 5q13.2

Domains: pkinase, TyrKc, S_TKc

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency, Transcription Factors





CDK7 (NM_001799) Human Tagged ORF Clone Lentiviral Particle - RC202736L2V

Protein Pathways: Cell cycle, Nucleotide excision repair

MW: 39.1 kDa

Gene Summary: The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK)

family. CDK family members are highly similar to the gene products of Saccharomyces cerevisiae cdc28, and Schizosaccharomyces pombe cdc2, and are known to be important regulators of cell cycle progression. This protein forms a trimeric complex with cyclin H and MAT1, which functions as a Cdk-activating kinase (CAK). It is an essential component of the transcription factor TFIIH, that is involved in transcription initiation and DNA repair. This protein is thought to serve as a direct link between the regulation of transcription and the cell

cycle. [provided by RefSeq, Jul 2008]