

## Product datasheet for **RC202743L3V**

### HLA-DRB4 (NM\_021983) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	HLA-DRB4 (NM_021983) Human Tagged ORF Clone Lentiviral Particle
Symbol:	HLA-DRB4
Synonyms:	DR4; DRB4; HLA-DR4B; HLA-DRB4*
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_021983
ORF Size:	798 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202743).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_021983.4</a> , <a href="#">NP_068818.4</a>
RefSeq Size:	1193 bp
RefSeq ORF:	801 bp
Locus ID:	3126
UniProt ID:	<a href="#">P13762</a>
Cytogenetics:	6p21.3
Domains:	MHC_II_beta, ig, IGc1
Protein Families:	Transmembrane



[View online »](#)

**Protein Pathways:** Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Hematopoietic cell lineage, Systemic lupus erythematosus, Type I diabetes mellitus, Viral myocarditis

**MW:** 29.9 kDa

**Gene Summary:** HLA-DRB4 belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells. The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. There are multiple pseudogenes of this gene. [provided by RefSeq, Feb 2020]