

## Product datasheet for RC202755L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Ribonuclease A (RNASE1) (NM 002933) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Ribonuclease A (RNASE1) (NM\_002933) Human Tagged ORF Clone Lentiviral Particle

Symbol: Ribonuclease A
Synonyms: RAC1; RIB1; RNS1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM\_002933

ORF Size: 468 bp

**ORF Nucleotide** 

TI 005

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC202755).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002933.3

 RefSeq Size:
 820 bp

 RefSeq ORF:
 471 bp

 Locus ID:
 6035

 UniProt ID:
 P07998

 Cytogenetics:
 14q11.2

 Domains:
 RNAse\_Pc

**Protein Families:** Secreted Protein, Transmembrane





## Ribonuclease A (RNASE1) (NM\_002933) Human Tagged ORF Clone Lentiviral Particle – RC202755L2V

**MW:** 17.6 kDa

**Gene Summary:** 

This gene encodes a member of the pancreatic-type of secretory ribonucleases, a subset of the ribonuclease A superfamily. The encoded endonuclease cleaves internal phosphodiester RNA bonds on the 3'-side of pyrimidine bases. It prefers poly(C) as a substrate and hydrolyzes 2',3'-cyclic nucleotides, with a pH optimum near 8.0. The encoded protein is monomeric and more commonly acts to degrade ds-RNA over ss-RNA. Alternative splicing occurs at this locus and four transcript variants encoding the same protein have been identified. [provided by RefSeq, Jul 2008]