

Product datasheet for **RC202799**

C4BPB (NM_001017365) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	C4BPB (NM_001017365) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	C4BPB
Synonyms:	C4BP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC202799 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGTTTTTTTGGTGTGCGTGCTGTCTTATGGTTGCGTGGCGAGTTCTGCTTCAGATGAGCACTGTCCAG
AGCTTCCTCCAGTGGACAATAGCATATTTGTCGCAAAGGAGGTGGAAGGACAGATTCTGGGGACTTACGT
TTGTATCAAGGGCTACCACCTGGTAGGAAAGAAGACCCTTTTTTGCAATGCCTCTAAGGAGTGGGATAAC
ACCACTACTGAGTGCCGCTTGGCCACTGTCCTGATCCTGTGCTGGTGAATGGAGAGTTCAGTTCTTCAG
GGCCTGTGAATGTAAGTGACAAAATCACGTTTATGTGCAATGACCACTACATCCTCAAGGGCAGCAATCG
GAGCCAGTGTCTAGAGGACCACACCTGGGCACCTCCCTTTCCCATCTGCAAAAGTAGGGACTGTGACCCT
CCTGGGAATCCAGTTCATGGCTATTTTGAAGGAAATAACTTCACCTTAGGATCCACCATTAGTTATTACT
GTGAAGACAGGTACTACTTAGTGGGCGTGCAGGAGCAGCAATGCGTTGATGGGGAGTGGAGCAGTGCAGT
TCCAGTCTGCAAGTTGATCCAGGAAGCTCCCAAACAGAGTGTGAGAAGGCACTTCTTGCTTTTCAGGAG
AGTAAGAACCTCTGCGAAGCCATGGAGAACTTTATGCAACAATTAAGGAAAGTGGCATGACAATGGAGG
AGCTAAAATATTCTCTGGAGCTGAAGAAAGCTGAGTTGAAGGCAAAATTGTTG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC202799 protein sequence
 Red=Cloning site Green=Tags(s)

MFFWCACCLMVAWRVSASDEHCPPELPPVDNSIFVAKEVEGQILGTYYVCIKGYHLVGKKTLCNASKEWDN
 TTTECRIGHCPDPVLVNGEFSSSGPVNVSDKITFMCNDHYILKGSNRSQCLEDHTWAPPFPICKSRDCDP
 PGNPVHGYFEGNFTLGSTISYYCEDRYLVGVQEQQCVDGEWSSALPVCKLIQEAPKPECEKALLAFQE
 SKNLCEAMENFMQQLKESGMTMEELKYSLELKKAEKAKLL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6628_c09.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001017365

ORF Size: 753 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001017365.3</u>
RefSeq Size:	968 bp
RefSeq ORF:	759 bp
Locus ID:	725
UniProt ID:	<u>P20851</u>
Cytogenetics:	1q32.1
Protein Pathways:	Complement and coagulation cascades
MW:	28.3 kDa
Gene Summary:	<p>This gene encodes a member of a superfamily of proteins composed predominantly of tandemly arrayed short consensus repeats of approximately 60 amino acids. A single, unique beta-chain encoded by this gene assembles with seven identical alpha-chains into the predominant isoform of C4b-binding protein, a multimeric protein that controls activation of the complement cascade through the classical pathway. C4b-binding protein has a regulatory role in the coagulation system also, mediated through the beta-chain binding of protein S, a vitamin K-dependent protein that serves as a cofactor of activated protein C. The genes encoding both alpha and beta chains are located adjacent to each other on human chromosome 1 in the regulator of complement activation gene cluster. Alternative splicing gives rise to multiple transcript variants. [provided by RefSeq, Jul 2008]</p>


Product images:



Circular map for RC202799



Western blot validation of overexpression lysate (Cat# [LY422641]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC219216] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified C4BPB protein (Cat# [TP302799]). The protein was produced from HEK293T cells transfected with C4BPB cDNA clone (Cat# RC202799) using MegaTran 2.0 (Cat# [TT210002]).