

## Product datasheet for RC202808L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Geminin (GMNN) (NM 015895) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Geminin (GMNN) (NM\_015895) Human Tagged ORF Clone Lentiviral Particle

Symbol: Geminin

**Synonyms:** Gem; MGORS6

Mammalian Cell

Selection:

ACCN:

None

NM 015895

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ORF Size: 627 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202808).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 015895.3

 RefSeq Size:
 1275 bp

 RefSeq ORF:
 630 bp

 Locus ID:
 51053

 UniProt ID:
 075496

 Cytogenetics:
 6p22.3

**Protein Families:** Druggable Genome, Stem cell - Pluripotency

MW: 23.6 kDa





## **Gene Summary:**

This gene encodes a protein that plays a critical role in cell cycle regulation. The encoded protein inhibits DNA replication by binding to DNA replication factor Cdt1, preventing the incorporation of minichromosome maintenance proteins into the pre-replication complex. The encoded protein is expressed during the S and G2 phases of the cell cycle and is degraded by the anaphase-promoting complex during the metaphase-anaphase transition. Increased expression of this gene may play a role in several malignancies including colon, rectal and breast cancer. Alternatively spliced transcript variants have been observed for this gene, and two pseudogenes of this gene are located on the short arm of chromosome 16. [provided by RefSeq, Oct 2011]