

## Product datasheet for **RC202878L1V**

### **DUSP2 (NM\_004418) Human Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	DUSP2 (NM_004418) Human Tagged ORF Clone Lentiviral Particle
Symbol:	DUSP2
Synonyms:	PAC-1; PAC1
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_004418
ORF Size:	942 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202878).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_004418.2</a>
RefSeq Size:	1708 bp
RefSeq ORF:	945 bp
Locus ID:	1844
UniProt ID:	<a href="#">Q05923</a>
Cytogenetics:	2q11.2
Domains:	DSPc, RHOD, PTPc_motif
Protein Families:	Druggable Genome, Phosphatase



[View online »](#)

**Protein Pathways:** MAPK signaling pathway

**MW:** 34.4 kDa

**Gene Summary:** The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1 and ERK2, is predominantly expressed in hematopoietic tissues, and is localized in the nucleus. [provided by RefSeq, Jul 2008]