

Product datasheet for RC202892L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HLA DOB (HLA-DOB) (NM_002120) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HLA DOB (HLA-DOB) (NM_002120) Human Tagged ORF Clone Lentiviral Particle

Symbol: HLA DOB

Synonyms: DOB; HLA_DOB

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 002120

ORF Size: 819 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202892).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002120.2</u>

 RefSeq Size:
 1388 bp

 RefSeq ORF:
 822 bp

 Locus ID:
 3112

 UniProt ID:
 P13765

 Cytogenetics:
 6p21.32

Protein Families: Transmembrane





HLA DOB (HLA-DOB) (NM_002120) Human Tagged ORF Clone Lentiviral Particle - RC202892L3V

Protein Pathways: Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid

disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Systemic lupus

erythematosus, Type I diabetes mellitus, Viral myocarditis

MW: 30.8 kDa

Gene Summary: HLA-DOB belongs to the HLA class II beta chain paralogues. This class II molecule is a

heterodimer consisting of an alpha (DOA) and a beta chain (DOB), both anchored in the membrane. It is located in intracellular vesicles. DO suppresses peptide loading of MHC class II molecules by inhibiting HLA-DM. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon

5 encodes the cytoplasmic tail. [provided by RefSeq, Jul 2008]