

Product datasheet for RC202896L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cystatin SN (CST1) (NM_001898) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cystatin SN (CST1) (NM 001898) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cystatin SN Puromycin

Mammalian Cell

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag:

ACCN: NM_001898

ORF Size: 423 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC202896).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001898.2, NP 001889.2

RefSeq Size: 782 bp RefSeq ORF: 426 bp Locus ID: 1469 **UniProt ID:** P01037

Cytogenetics: 20p11.21

Domains: CY

MW: 16.4 kDa







Gene Summary:

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a cysteine proteinase inhibitor found in saliva, tears, urine, and seminal fluid. [provided by RefSeq, Jul 2008]