

Product datasheet for RC203089L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RNF2 (NM_007212) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RNF2 (NM_007212) Human Tagged ORF Clone Lentiviral Particle

Symbol: RNF2

Synonyms: BAP-1; BAP1; DING; HIPI3; RING1B; RING2

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_007212

 ORF Size:
 1008 bp

ORF Nucleotide

OTI Disclaimer:

1000 pp

Sequence:

The ORF insert of this clone is exactly the same as(RC203089).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 007212.3

 RefSeq Size:
 3551 bp

 RefSeq ORF:
 1011 bp

 Locus ID:
 6045

 UniProt ID:
 Q99496

 Cytogenetics:
 1q25.3

Domains: RING

Protein Families: Druggable Genome, Transcription Factors







MW: 37.7 kDa

Gene Summary:

Polycomb group (PcG) of proteins form the multiprotein complexes that are important for the transcription repression of various genes involved in development and cell proliferation. The protein encoded by this gene is one of the PcG proteins. It has been shown to interact with, and suppress the activity of, transcription factor CP2 (TFCP2/CP2). Studies of the mouse counterpart suggested the involvement of this gene in the specification of anterior-posterior axis, as well as in cell proliferation in early development. This protein was also found to interact with huntingtin interacting protein 2 (HIP2), an ubiquitin-conjugating enzyme, and possess ubiquitin ligase activity. [provided by RefSeq, Jul 2008]