

## Product datasheet for RC203102L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BCA1 (CXCL13) (NM\_006419) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: BCA1 (CXCL13) (NM 006419) Human Tagged ORF Clone Lentiviral Particle

Symbol: BCA1

Synonyms: ANGIE; ANGIE2; BCA-1; BCA1; BLC; BLR1L; SCYB13

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 006419

ORF Size: 327 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203102).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 006419.1, NP 006410.1

 RefSeq Size:
 1219 bp

 RefSeq ORF:
 330 bp

 Locus ID:
 10563

 UniProt ID:
 043927

 Cytogenetics:
 4q21.1

Domains: IL8

**Protein Families:** Druggable Genome, Secreted Protein





## BCA1 (CXCL13) (NM\_006419) Human Tagged ORF Clone Lentiviral Particle - RC203102L3V

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

**MW:** 12.7 kDa

**Gene Summary:** B lymphocyte chemoattractant, independently cloned and named Angie, is an antimicrobial

peptide and CXC chemokine strongly expressed in the follicles of the spleen, lymph nodes, and Peyer's patches. It preferentially promotes the migration of B lymphocytes (compared to T cells and macrophages), apparently by stimulating calcium influx into, and chemotaxis of, cells expressing Burkitt's lymphoma receptor 1 (BLR-1). It may therefore function in the

homing of B lymphocytes to follicles. [provided by RefSeq, Oct 2014]