

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC203115L1V

Cystatin A (CSTA) (NM_005213) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Cystatin A (CSTA) (NM_005213) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Cystatin A
Synonyms:	AREI; PSS4; STF1; STFA
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_005213
ORF Size:	294 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203115).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 005213.3</u>
RefSeq Size:	838 bp
RefSeq ORF:	297 bp
Locus ID:	1475
UniProt ID:	<u>P01040</u>
Cytogenetics:	3q21.1
Domains:	CY
MW:	11 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Cystatin A (CSTA) (NM_005213) Human Tagged ORF Clone Lentiviral Particle – RC203115L1V Cystatin A (CSTA) (NM_005213) Human Tagged ORF Clone Lentiviral Particle – RC203115L1V

Gene Summary:The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences.
Some of the members are active cysteine protease inhibitors, while others have lost or
perhaps never acquired this inhibitory activity. There are three inhibitory families in the
superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This
gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes
with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of
cornified cell envelope in keratinocytes and plays a role in epidermal development and
maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer.
[provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US