

## Product datasheet for RC203131L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RPL35A (NM\_000996) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: RPL35A (NM 000996) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPL35A

Synonyms: DBA5; eL33; L35A

**Mammalian Cell** 

Puromycin

Selection:

ACCN:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 000996

Tag: Myc-DDK

ORF Size: 330 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203131).

Sequence:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000996.2

 RefSeq Size:
 511 bp

 RefSeq ORF:
 333 bp

 Locus ID:
 6165

 UniProt ID:
 P18077

 Cytogenetics:
 3q29

**Domains:** Ribosomal\_L35Ae

**Protein Pathways:** Ribosome





ORIGENE

**MW:** 12.5 kDa

**Gene Summary:** 

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L35AE family of ribosomal proteins. It is located in the cytoplasm. The rat protein has been shown to bind to both initiator and elongator tRNAs, and thus, it is located at the P site, or P and A sites, of the ribosome. Although this gene was originally mapped to chromosome 18, it has been established that it is located at 3q29-qter. Alternative splicing results in multiple transcript variants. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Oct 2015]