

Product datasheet for **RC203142L2V**

Tissue Factor (F3) (NM_001993) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Tissue Factor (F3) (NM_001993) Human Tagged ORF Clone Lentiviral Particle
Symbol:	F3
Synonyms:	CD142; TF; TFA
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_001993
ORF Size:	885 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203142).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001993.2
RefSeq Size:	2393 bp
RefSeq ORF:	888 bp
Locus ID:	2152
UniProt ID:	P13726
Cytogenetics:	1p21.3
Domains:	Tissue_fac
Protein Families:	Druggable Genome, Transmembrane



[View online »](#)

Protein Pathways: Complement and coagulation cascades

MW: 33.1 kDa

Gene Summary: This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces, for example, on monocytes. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. Platelets and monocytes have been shown to express this coagulation factor under procoagulatory and proinflammatory stimuli, and a major role in HIV-associated coagulopathy has been described. Platelet-dependent monocyte expression of coagulation factor III has been described to be associated with Coronavirus Disease 2019 (COVID-19) severity and mortality. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Aug 2020]