

Product datasheet for RC203143L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cathepsin L (CTSL) (NM_001912) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cathepsin L (CTSL) (NM_001912) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cathepsin L

Synonyms: CATL; CTSL1; MEP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 001912

ORF Size: 999 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC203143).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001912.3

 RefSeq Size:
 1730 bp

 RefSeq ORF:
 1002 bp

 Locus ID:
 1514

 UniProt ID:
 P07711

 Cytogenetics:
 9q21.33

Domains: Pept_C1

Protein Families: Druggable Genome, Protease





Cathepsin L (CTSL) (NM_001912) Human Tagged ORF Clone Lentiviral Particle - RC203143L3V

Protein Pathways: Antigen processing and presentation, Lysosome

MW: 37.5 kDa

Gene Summary: The protein encoded by this gene is a lysosomal cysteine proteinase that plays a major role in

intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. The encoded protein has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. This protein, which is a member of the peptidase C1 family, is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Additionally, this protein cleaves the S1 subunit of the SARS-CoV-2 spike protein, which is necessary for

entry of the virus into the cell. [provided by RefSeq, Aug 2020]