

# Product datasheet for RC203144L1

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Corticotropin Releasing Factor (CRH) (NM\_000756) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Corticotropin Releasing Factor (CRH) (NM\_000756) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Corticotropin Releasing Factor

Synonyms: CRF; CRH1

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC203144).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_000756

ORF Size: 588 bp



### Corticotropin Releasing Factor (CRH) (NM\_000756) Human Tagged Lenti ORF Clone - RC203144L1

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 000756.1</u>

**RefSeq Size:** 1434 bp

RefSeg ORF: 591 bp

**Locus ID:** 1392

UniProt ID: P06850

Cytogenetics: 8q13.1

Domains: CRF

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Long-term depression


MW: 21.4 kDa



#### **Gene Summary:**


This gene encodes a member of the corticotropin-releasing factor family. The encoded preproprotein is proteolytically processed to generate the mature neuropeptide hormone. In response to stress, this hormone is secreted by the paraventricular nucleus (PVN) of the hypothalamus, binds to corticotropin releasing hormone receptors and stimulates the release of adrenocorticotropic hormone from the pituitary gland. Marked reduction in this protein has been observed in association with Alzheimer's disease. Autosomal recessive hypothalamic corticotropin deficiency has multiple and potentially fatal metabolic consequences including hypoglycemia and hepatitis. In addition to production in the hypothalamus, this protein is also synthesized in peripheral tissues, such as T lymphocytes, and is highly expressed in the placenta. In the placenta it is a marker that determines the length of gestation and the timing of parturition and delivery. A rapid increase in circulating levels of the hormone occurs at the onset of parturition, suggesting that, in addition to its metabolic functions, this protein may act as a trigger for parturition. [provided by RefSeq, Nov 2015]

# **Product images:**



Circular map for RC203144L1





Double digestion of RC203144L1 using Sgfl and Mlul