

Product datasheet for RC203155L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NDUFB6 (NM_002493) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: NDUFB6 (NM_002493) Human Tagged ORF Clone Lentiviral Particle

Symbol: NDUFB6
Synonyms: B17; CI

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002493

ORF Size: 384 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC203155).

•

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002493.3

 RefSeq Size:
 873 bp

 RefSeq ORF:
 387 bp

 Locus ID:
 4712

 UniProt ID:
 095139

 Cytogenetics:
 9p21.1

Protein Families: Transmembrane





NDUFB6 (NM_002493) Human Tagged ORF Clone Lentiviral Particle - RC203155L4V

Protein Pathways: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation,

Parkinson's disease

MW: 15.5 kDa

Gene Summary: The protein encoded by this gene is a subunit of the multisubunit NADH:ubiquinone

oxidoreductase (complex I). Mammalian complex I is composed of 45 different subunits. It locates at the mitochondrial inner membrane. This protein has NADH dehydrogenase activity and oxidoreductase activity. It transfers electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone. Alternative splicing

occurs at this locus and three transcript variants encoding distinct isoforms have been

identified. [provided by RefSeq, Jan 2011]