

## Product datasheet for **RC203163L3V**

### SOCS2 (NM\_003877) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	SOCS2 (NM_003877) Human Tagged ORF Clone Lentiviral Particle
Symbol:	SOCS2
Synonyms:	CIS2; Cish2; SOCS-2; SSI-2; SSI2; STAT12
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_003877
ORF Size:	594 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203163).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_003877.3</a>
RefSeq Size:	2759 bp
RefSeq ORF:	597 bp
Locus ID:	8835
UniProt ID:	<a href="#">O14508</a>
Cytogenetics:	12q22
Domains:	SH2, SOCS
Protein Families:	Druggable Genome



[View online »](#)

---

<b>Protein Pathways:</b>	Insulin signaling pathway, Jak-STAT signaling pathway, Type II diabetes mellitus
<b>MW:</b>	22.2 kDa
<b>Gene Summary:</b>	<p>This gene encodes a member of the suppressor of cytokine signaling (SOCS) family. SOCS family members are cytokine-inducible negative regulators of cytokine receptor signaling via the Janus kinase/signal transducer and activation of transcription pathway (the JAK/STAT pathway). SOCS family proteins interact with major molecules of signaling complexes to block further signal transduction, in part, by proteasomal depletion of receptors or signal-transducing proteins via ubiquitination. The expression of this gene can be induced by a subset of cytokines, including erythropoietin, GM-CSF, IL10, interferon (IFN)-gamma and by cytokine receptors such as growth hormone receptor. The protein encoded by this gene interacts with the cytoplasmic domain of insulin-like growth factor-1 receptor (IGF1R) and is thought to be involved in the regulation of IGF1R mediated cell signaling. This gene has pseudogenes on chromosomes 20 and 22. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2012]</p>