

## Product datasheet for RC203176L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MPZL (MPZL1) (NM\_003953) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: MPZL (MPZL1) (NM 003953) Human Tagged ORF Clone Lentiviral Particle

Symbol: MPZL

**Synonyms:** MPZL1b; PZR; PZR1b; PZRa; PZRb

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_003953

ORF Size: 807 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203176).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 003953.4

 RefSeq Size:
 5026 bp

 RefSeq ORF:
 810 bp

 Locus ID:
 9019

 UniProt ID:
 095297

 Cytogenetics:
 1q24.2

Domains: IGv, IG

**Protein Families:** Druggable Genome, Transmembrane





## MPZL (MPZL1) (NM\_003953) Human Tagged ORF Clone Lentiviral Particle - RC203176L2V

**Protein Pathways:** Cell adhesion molecules (CAMs)

MW: 29.1 kDa

Gene Summary: Cell surface receptor, which is involved in signal transduction processes. Recruits

PTPN11/SHP-2 to the cell membrane and is a putative substrate of PTPN11/SHP-2. Is a major receptor for concanavalin-A (ConA) and is involved in cellular signaling induced by ConA, which probably includes Src family tyrosine-protein kinases. Isoform 3 seems to have a dominant negative role; it blocks tyrosine phosphorylation of MPZL1 induced by ConA. Isoform 1, but not isoform 2 and isoform 3, may be involved in regulation of integrin-

mediated cell motility.[UniProtKB/Swiss-Prot Function]