

## Product datasheet for RC203181L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MARCKS like protein (MARCKSL1) (NM\_023009) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** MARCKS like protein (MARCKSL1) (NM\_023009) Human Tagged ORF Clone Lentiviral Particle

Symbol: MARCKS like protein

Synonyms: F52; MACMARCKS; MLP; MLP1; MRP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_023009

ORF Size: 585 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203181).

Sequence:

Domains:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 023009.4

 RefSeq Size:
 1562 bp

 RefSeq ORF:
 588 bp

 Locus ID:
 65108

 UniProt ID:
 P49006

 Cytogenetics:
 1p35.1

**Protein Pathways:** Fc gamma R-mediated phagocytosis

**MARCKS** 





MARCKS like protein (MARCKSL1) (NM\_023009) Human Tagged ORF Clone Lentiviral Particle – RC203181L4V

**MW:** 19.5 kDa

**Gene Summary:** This gene encodes a member of the myristoylated alanine-rich C-kinase substrate (MARCKS)

family. Members of this family play a role in cytoskeletal regulation, protein kinase C signaling and calmodulin signaling. The encoded protein affects the formation of adherens junction. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are located on the long arm of chromosomes 6 and 10. [provided by RefSeq, Jun 2012]