

Product datasheet for RC203182L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SDHB (NM_003000) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SDHB (NM 003000) Human Tagged ORF Clone Lentiviral Particle

Symbol: SDHB

Synonyms: CWS2; IP; MC2DN4; PGL4; SDH; SDH1; SDH2; SDHIP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 003000

ORF Size: 840 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC203182).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003000.2</u>

 RefSeq Size:
 1161 bp

 RefSeq ORF:
 843 bp

 Locus ID:
 6390

 UniProt ID:
 P21912

 Cytogenetics:
 1p36.13

Domains: fer2

Protein Families: Druggable Genome





SDHB (NM_003000) Human Tagged ORF Clone Lentiviral Particle - RC203182L3V

Protein Pathways: Alzheimer's disease, Citrate cycle (TCA cycle), Huntington's disease, Metabolic pathways,

Oxidative phosphorylation, Parkinson's disease

MW: 31.6 kDa

Gene Summary: Complex II of the respiratory chain, which is specifically involved in the oxidation of succinate,

carries electrons from FADH to CoQ. The complex is composed of four nuclear-encoded subunits and is localized in the mitochondrial inner membrane. The iron-sulfur subunit is highly conserved and contains three cysteine-rich clusters which may comprise the iron-sulfur centers of the enzyme. Sporadic and familial mutations in this gene result in paragangliomas and pheochromocytoma, and support a link between mitochondrial

dysfunction and tumorigenesis. [provided by RefSeq, Jul 2008]