

Product datasheet for **RC203276**

Tropomyosin 3 (TPM3) (NM_152263) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Tag:	Myc-DDK
Symbol:	Tropomyosin 3
Synonyms:	CAPM1; CFTD; HEL-189; HEL-S-82p; hscp30; NEM1; OK/SW-cl.5; TM-5; TM3; TM5; TM30; TM30nm; TPM3nu; TPMsk3; TRK
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC203276 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGC**

ATGATGGAGGCCATCAAGAAAAAGATGCAGATGCTGAAGTTAGACAAGGAGAATGCTCTGGATCGGGCAG
AGCAAGCTGAAGCTGAGCAGAAGCAGGCAGAAGAAAGTAACAGCTGGAGGATGAGCTGGCAGCCAT
GCAGAAGAAGCTGAAAGGGACAGAGGATGAGCTGGACAAGTATTCTGAAGCTTTGAAGGATGCCAGGAG
AAGCTGGAAGCTGGCAGAGAAGAAGGCTGCTGATGCTGAGGCTGAGGTGGCCTCCTTGAACCGTAGGATCC
AGCTGGTTGAAGAAGAGCTGGACCGTCTCAGGAGCGCCTGGCCACTGCCCTGCAAAAGCTGGAAGAAGC
TGAAAAAGCTGCTGATGAGAGTGAGAGAGGTATGAAGTTATTGAAAACCGGCCCTTAAAGATGAAGAA
AAGATGGAAGCTCCAGGAAATCCAACCTCAAAGAAGCTAAGCACATTGCAGAAGAGGCAGATAGGAAGTATG
AAGAGGTGGCTCGTAAGTTGGTGATCATTGAAGGAGACTTGGAACGCACAGAGGAACGAGCTGAGCTGGC
AGAGTCTAAGTGTTCTGAGCTGGAGGAGGAGCTGAAGAATGTACCAACAACCTCAAGTCTCTTGAGGCT
CAGGCGGAGAAGTACTCTCAAAAAGAAGATAAATATGAGGAAGAAATCAAGATTCTTACTGATAAACTCA
AGGAGGCAGAGACCCGTGCTGAGTTTGCTGAGAGATCGGTAGCCAAGCTGGAAAAGACAATTGATGACCT
GGAAGATGAGCTCTATGCCCAGAACTGAAGTACAAGGCCATTAGCGAGGAGCTGGACCACGCCCTCAAT
GACATGACCTCTATA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC203276 protein sequence
 Red=Cloning site Green=Tags(s)


MMEAIKKKMQLKLDKENALDRAEQAEAEQQAEEERSKQLEDELAAMQKKLKGTEDELDKYSEALKDAQE
 KLELAEKKAADAEVASLNRRIQLVVEELDRAQERLATALQKLEEAKAADESERGMKVIENTRALKDEE
 KMELQEIQLKEAKHIAEEADRKYEEVARKLVIIEGDLERTEERAELAESKCSLEEELKNVTNNLKSLEA
 QAEKYSQKEDKYEIEIKILTDKLKEATRAEFAERSVAKLEKTIDDLDELDYQKLKYKAISEELDHALN
 DMTSI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6416_b07.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_152263

ORF Size: 855 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_152263.4](#)

RefSeq Size: 7109 bp

RefSeq ORF: 858 bp

Locus ID: 7170

UniProt ID: [P06753](#)

Cytogenetics: 1q21.3


Domains: Tropomyosin

Protein Pathways: Cardiac muscle contraction, Dilated cardiomyopathy, Hypertrophic cardiomyopathy (HCM), Pathways in cancer, Thyroid cancer


MW: 32.9 kDa

Gene Summary: This gene encodes a member of the tropomyosin family of actin-binding proteins. Tropomyosins are dimers of coiled-coil proteins that provide stability to actin filaments and regulate access of other actin-binding proteins. Mutations in this gene result in autosomal dominant nemaline myopathy and other muscle disorders. This locus is involved in translocations with other loci, including anaplastic lymphoma receptor tyrosine kinase (ALK) and neurotrophic tyrosine kinase receptor type 1 (NTRK1), which result in the formation of fusion proteins that act as oncogenes. There are numerous pseudogenes for this gene on different chromosomes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2013]


Product images:



Circular map for RC203276



Western blot validation of overexpression lysate (Cat# [LY407686]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC203276 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified TPM3 protein (Cat# [TP303276]). The protein was produced from HEK293T cells transfected with TPM3 cDNA clone (Cat# RC203276) using MegaTran 2.0 (Cat# [TT210002]).