

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC203435L1V

## HIP55 (DBNL) (NM\_014063) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	HIP55 (DBNL) (NM_014063) Human Tagged ORF Clone Lentiviral Particle
Symbol:	HIP55
Synonyms:	ABP1; HIP-55; HIP55; SH3P7
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_014063
ORF Size:	1293 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203435).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 014063.5</u>
RefSeq Size:	2213 bp
RefSeq ORF:	1296 bp
Locus ID:	28988
UniProt ID:	<u>Q9UJU6</u>
Cytogenetics:	7p13
Domains:	SH3, ADF
MW:	48.3 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Adapter protein that binds F-actin and DNM1, and thereby plays a role in receptor-mediated<br/>endocytosis. Plays a role in the reorganization of the actin cytoskeleton, formation of cell<br/>projections, such as neurites, in neuron morphogenesis and synapse formation via its<br/>interaction with WASL and COBL. Does not bind G-actin and promote actin polymerization by<br/>itself. Required for the formation of organized podosome rosettes (By similarity). May act as a<br/>common effector of antigen receptor-signaling pathways in leukocytes. Acts as a key<br/>component of the immunological synapse that regulates T-cell activation by bridging TCRs<br/>and the actin cytoskeleton to gene activation and endocytic processes.[UniProtKB/Swiss-Prot<br/>Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US