

Product datasheet for RC203461L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SHMT1 (NM 004169) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SHMT1 (NM 004169) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CSHMT; SHMT Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag: NM 004169 ACCN:

ORF Size: 1449 bp

ORF Nucleotide

Sequence:

UniProt ID:

The ORF insert of this clone is exactly the same as(RC203461).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004169.3

P34896

RefSeq Size: 2553 bp RefSeq ORF: 1452 bp

Locus ID: 6470

Cytogenetics: 17p11.2

Domains: SHMT





SHMT1 (NM_004169) Human Tagged ORF Clone Lentiviral Particle - RC203461L3V

Protein Pathways: Cyanoamino acid metabolism, Glycine, serine and threonine metabolism, Metabolic

pathways, Methane metabolism, One carbon pool by folate

MW: 53.1 kDa

Gene Summary: This gene encodes the cytosolic form of serine hydroxymethyltransferase, a pyridoxal

phosphate-containing enzyme that catalyzes the reversible conversion of serine and tetrahydrofolate to glycine and 5,10-methylene tetrahydrofolate. This reaction provides one-carbon units for synthesis of methionine, thymidylate, and purines in the cytoplasm. This

gene is located within the Smith-Magenis syndrome region on chromosome 17. A

pseudogene of this gene is located on the short arm of chromosome 1. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Aug 2013]