

## Product datasheet for **RC203480**

### Phosphatidic acid phosphatase type 2B (PLPP3) (NM\_003713) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Phosphatidic acid phosphatase type 2B (PLPP3) (NM_003713) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Phosphatidic acid phosphatase type 2B
Synonyms:	Dri42; LPP3; PAP2B; PPAP2B; VCIP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC203480 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGCAAACTACAAGTACGACAAAGCGATCGTCCCGGAGAGCAAGAACGGCGGCAGCCCGGCGCTCAACA  
ACAACCCGAGGAGGAGCGGCAGCAAGCGGGTGTGCTCATCTGCCTCGACCTCTTCTGCCTTTCATGGC  
GGGCTCCCTTCTCATCATCGAGACAAGCACCATCAAGCCTTACCACCGAGGGTTTTACTGCAATGAT  
GAGAGCATCAAGTACCCACTGAAAAGTGGTGGAGACAATAATGACGCTGTGCTCTGTGCCGTGGGGATCG  
TCATTGCCATCCTCGCGATCATCACGGGGGAATTCACCGGATCTATTACCTGAAGAAGTCGCGGTGCGAC  
GATTCAGAACCCCTACGTGGCAGCACTCTATAAGCAAGTGGGCTGCTTCCTCTTTGGCTGTGCCATCAGC  
CAGTCTTTCACAGACATTGCCAAAGTGTCCATAGGGCGCCTGCGTCCCTCACTTCTTGAGTGTCTGCAACC  
CTGATTTTCAGCCAGATCAACTGCTCTGAAGGCTACATTCAGAAGTACAGATGCAGAGGTGATGACAGCAA  
AGTCCAGGAAGCCAGGAAGTCTTCTTCTGCGCATGCCTCCTTCTCCATGTACACTATGCTGTATTTG  
GTGCTATACCTGCAGGCCCGCTTCACTTGGCGAGGAGCCCGCCTGCTCCGGCCCCCTCGCAGTTCACTT  
TGATCATGATGGCCTTCTACACGGGACTGTCTCGCGTATCAGACCACAAGCACCATCCAGTGATGTTCT  
GGCAGGATTTGCTCAAGGAGCCCTGGTGGCCTGCTGCATAGTTTTCTTCTGCTGTGACCTCTTCAAGACT  
AAGATGACGCTCTCCCTGCTGCCCTGCTATCCGGAAGGAAATCCTTTCACCTGTGGACATTATTGACA  
GGAACAATCACCAACATGATG

**ACGCGT**ACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >RC203480 protein sequence  
Red=Cloning site Green=Tags(s)

MQNYKYDKAIVPESKNGGSPALNNPNRRSGSKRVLLICLDLFCLFMAGLPFLIIETSTIKPYHRGFYCNDESIKYPLKTGETINDAVLCAVGIVIAILAIITGEFYRIYLLKSRSTIQNPYVAALYKQVGCFLFGCAISQSFTDIKVSIGRLRPHFLSVCNPDFSQINCSEGYIQNYRCRGDDSKVQEARKSFFSGHASF SMYTMLYLVLYLQARFTWRGARLLRPLLQFTLIMMAFYTGLSRVSDHKHHPSDVLAGFAQGALVACCI VFFVSDLFKTKMTLSLPAPAIRKEILSPVDIIDRNHHNMM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:** [https://cdn.origene.com/chromatograms/mk6424\\_e09.zip](https://cdn.origene.com/chromatograms/mk6424_e09.zip)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_003713

**ORF Size:** 933 bp


**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_003713.2</a>
<b>RefSeq Size:</b>	3324 bp
<b>RefSeq ORF:</b>	936 bp
<b>Locus ID:</b>	8613
<b>UniProt ID:</b>	<a href="#">O14495</a>
<b>Cytogenetics:</b>	1p32.2
<b>Domains:</b>	acidPPc
<b>Protein Families:</b>	Druggable Genome, Transmembrane
<b>Protein Pathways:</b>	Ether lipid metabolism, Fc gamma R-mediated phagocytosis, Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways, Sphingolipid metabolism
<b>MW:</b>	35.1 kDa
<b>Gene Summary:</b>	The protein encoded by this gene is a member of the phosphatidic acid phosphatase (PAP) family. PAPs convert phosphatidic acid to diacylglycerol, and function in de novo synthesis of glycerolipids as well as in receptor-activated signal transduction mediated by phospholipase D. This protein is a membrane glycoprotein localized at the cell plasma membrane. It has been shown to actively hydrolyze extracellular lysophosphatidic acid and short-chain phosphatidic acid. The expression of this gene is found to be enhanced by epidermal growth factor in Hela cells. [provided by RefSeq, Mar 2010]


Product images:



Circular map for RC203480



Western blot validation of overexpression lysate (Cat# [LY406156]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC224959] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified PLPP3 protein (Cat# [TP303480]). The protein was produced from HEK293T cells transfected with PLPP3 cDNA clone (Cat# RC203480) using MegaTran 2.0 (Cat# [TT210002]).