

Product datasheet for RC203480L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phosphatidic acid phosphatase type 2B (PLPP3) (NM_003713) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Phosphatidic acid phosphatase type 2B (PLPP3) (NM_003713) Human Tagged ORF Clone

Lentiviral Particle

Symbol: Phosphatidic acid phosphatase type 2B

Synonyms: Dri42; LPP3; PAP2B; PPAP2B; VCIP

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 003713

ORF Size: 933 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC203480).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003713.3</u>

 RefSeq Size:
 3324 bp

 RefSeq ORF:
 936 bp

 Locus ID:
 8613

 UniProt ID:
 014495

 Cytogenetics:
 1p32.2

Domains: acidPPc





Phosphatidic acid phosphatase type 2B (PLPP3) (NM_003713) Human Tagged ORF Clone Lentiviral Particle – RC203480L1V

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Ether lipid metabolism, Fc gamma R-mediated phagocytosis, Glycerolipid metabolism,

Glycerophospholipid metabolism, Metabolic pathways, Sphingolipid metabolism

MW: 35.1 kDa

Gene Summary: The protein encoded by this gene is a member of the phosphatidic acid phosphatase (PAP)

family. PAPs convert phosphatidic acid to diacylglycerol, and function in de novo synthesis of glycerolipids as well as in receptor-activated signal transduction mediated by phospholipase D. This protein is a membrane glycoprotein localized at the cell plasma membrane. It has been shown to actively hydrolyze extracellular lysophosphatidic acid and short-chain

phosphatidic acid. The expression of this gene is found to be enhanced by epidermal growth

factor in Hela cells. [provided by RefSeq, Mar 2010]