

## Product datasheet for RC203540L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## B MyB (MYBL2) (NM 002466) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: B MyB (MYBL2) (NM 002466) Human Tagged ORF Clone Lentiviral Particle

Symbol: B MyB

Synonyms: B-MYB; BMYB

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_002466 **ORF Size:** 2100 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203540).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002466.2, NP 002457.1

 RefSeq Size:
 2785 bp

 RefSeq ORF:
 2103 bp

 Locus ID:
 4605

 UniProt ID:
 P10244

Cytogenetics: 20q13.12

**Domains:** myb\_DNA-binding

**Protein Families:** Druggable Genome, Stem cell - Pluripotency, Transcription Factors





**MW:** 78.8 kDa

**Gene Summary:** 

The protein encoded by this gene, a member of the MYB family of transcription factor genes, is a nuclear protein involved in cell cycle progression. The encoded protein is phosphorylated by cyclin A/cyclin-dependent kinase 2 during the S-phase of the cell cycle and possesses both activator and repressor activities. It has been shown to activate the cell division cycle 2, cyclin D1, and insulin-like growth factor-binding protein 5 genes. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2013]