

## Product datasheet for RC203585L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cofilin 1 (CFL1) (NM 005507) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Cofilin 1 (CFL1) (NM\_005507) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CFL; cofilin; HEL-S-15 Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 005507

**ORF Size:** 498 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203585).

Sequence: OTI Disclaimer:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 005507.2

RefSeq Size: 1260 bp RefSeq ORF: 501 bp Locus ID: 1072 **UniProt ID:** P23528 Cytogenetics: 11q13.1

**Protein Families:** Druggable Genome

ADF





## Cofilin 1 (CFL1) (NM\_005507) Human Tagged ORF Clone Lentiviral Particle - RC203585L1V

**Protein Pathways:** Axon guidance, Fc gamma R-mediated phagocytosis, Regulation of actin cytoskeleton

**MW:** 18.5 kDa

**Gene Summary:** The protein encoded by this gene can polymerize and depolymerize F-actin and G-actin in a

pH-dependent manner. Increased phosphorylation of this protein by LIM kinase aids in Rho-induced reorganization of the actin cytoskeleton. Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner. It is involved in the translocation of actin-cofilin complex from cytoplasm to nucleus.[supplied by OMIM, Apr

2004]