

Product datasheet for RC203616L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RPL29 (NM_000992) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RPL29 (NM 000992) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPL29

Synonyms: HIP; HUMRPL29; L29; RPL29P10; RPL29_3_370

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 000992

ORF Size: 477 bp

ORF Nucleotide

The OP

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC203616).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000992.2</u>

 RefSeq Size:
 737 bp

 RefSeq ORF:
 480 bp

 Locus ID:
 6159

 UniProt ID:
 P47914

 Cytogenetics:
 3p21.2

Domains: Ribosomal_L29e

Protein Pathways: Ribosome





ORIGENE

MW: 17.8 kDa

Gene Summary:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 60S subunit. The protein belongs to the L29E family of ribosomal proteins. The protein is also a peripheral membrane protein expressed on the cell surface that directly binds heparin. Although this gene was previously reported to map to 3q29-qter, it is believed that it is located at 3p21.3-p21.2. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]