

Product datasheet for **RC203652L1V**

ATP6V0C (NM_001694) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ATP6V0C (NM_001694) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ATP6V0C
Synonyms:	ATP6C; ATP6L; ATPL; VATL; Vma3; VPPC
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_001694
ORF Size:	465 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203652).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001694.2
RefSeq Size:	1180 bp
RefSeq ORF:	468 bp
Locus ID:	527
UniProt ID:	P27449
Cytogenetics:	16p13.3
Domains:	ATP-synt_C
Protein Families:	Transmembrane



[View online »](#)

Protein Pathways:	Epithelial cell signaling in Helicobacter pylori infection, Lysosome, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection
MW:	15.7 kDa
Gene Summary:	<p>This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c", and d. This gene encodes the V0 subunit c. Alternative splicing results in transcript variants. Pseudogenes have been identified on chromosomes 6 and 17. [provided by RefSeq, Nov 2010]</p>