

Product datasheet for RC203761L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VPAC2 (VIPR2) (NM_003382) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: VPAC2 (VIPR2) (NM_003382) Human Tagged ORF Clone Lentiviral Particle

Symbol: VPAC2

Synonyms: C16DUPq36.3; DUP7q36.3; PACAP-R-3; VIP-R-2; VPAC2; VPAC2R; VPCAP2R

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_003382 **ORF Size:** 1314 bp

ORF Nucleotide

131100

Sequence:

The ORF insert of this clone is exactly the same as(RC203761).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally accurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003382.3

 RefSeq Size:
 3961 bp

 RefSeq ORF:
 1317 bp

 Locus ID:
 7434

 UniProt ID:
 P41587

 Cytogenetics:
 7q36.3

Domains: 7tm_2, HormR

Protein Families: Druggable Genome, GPCR, Transmembrane





VPAC2 (VIPR2) (NM_003382) Human Tagged ORF Clone Lentiviral Particle - RC203761L2V

Protein Pathways: Neuroactive ligand-receptor interaction

MW: 49.5 kDa

Gene Summary: This gene encodes a receptor for vasoactive intestinal peptide, a small neuropeptide.

Vasoactive intestinal peptide is involved in smooth muscle relaxation, exocrine and endocrine secretion, and water and ion flux in lung and intestinal epithelia. Its actions are effected through integral membrane receptors associated with a guanine nucleotide binding protein

which activates adenylate cyclase. [provided by RefSeq, Aug 2011]