

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC203782L3V

## MLK3 (MAP3K11) (NM\_002419) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	MLK3 (MAP3K11) (NM_002419) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MLK3
Synonyms:	MEKK11; MLK-3; MLK3; PTK1; SPRK
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_002419
ORF Size:	2541 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203782).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 002419.3</u>
RefSeq Size:	3574 bp
RefSeq ORF:	2544 bp
Locus ID:	4296
UniProt ID:	<u>Q16584</u>
Cytogenetics:	11q13.1
Domains:	pkinase, TyrKc, SH3, S_TKc
Protein Families:	Druggable Genome, Protein Kinase



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	MLK3 (MAP3K11) (NM_002419) Human Tagged ORF Clone Lentiviral Particle – RC203782L3V
Protein Pathway	s: MAPK signaling pathway
MW:	92.5 kDa
Gene Summary:	The protein encoded by this gene is a member of the serine/threonine kinase family. This kinase contains a SH3 domain and a leucine zipper-basic motif. This kinase preferentially activates MAPK8/JNK kinase, and functions as a positive regulator of JNK signaling pathway. This kinase can directly phosphorylate, and activates IkappaB kinase alpha and beta, and is found to be involved in the transcription activity of NF-kappaB mediated by Rho family GTPases and CDC42. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US