

## Product datasheet for RC203789L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MCP4 (CCL13) (NM\_005408) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** MCP4 (CCL13) (NM\_005408) Human Tagged ORF Clone Lentiviral Particle

Symbol: MCP4

Synonyms: CKb10; MCP-4; NCC-1; NCC1; SCYA13; SCYL1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_005408

ORF Size: 294 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203789).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 005408.2

RefSeq Size: 861 bp
RefSeq ORF: 297 bp
Locus ID: 6357
UniProt ID: Q99616
Cytogenetics: 17q12

**Protein Families:** Druggable Genome, Secreted Protein





## MCP4 (CCL13) (NM\_005408) Human Tagged ORF Clone Lentiviral Particle - RC203789L4V

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction, NOD-like receptor

signaling pathway

MW: 11 kDa

**Gene Summary:** This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm

of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during artherosclerosis. [provided by RefSeq, Sep 2014]