

## **Product datasheet for RC203804**

## RGS13 (NM 144766) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** RGS13 (NM\_144766) Human Tagged ORF Clone

Tag:Myc-DDKSymbol:RGS13

Mammalian Cell Neomycin

Selection:

**Vector:** pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >RC203804 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGAGCAGGCGGAATTGTTGGATTTGTAAGATGTGCAGAGATGAATCTAAGAGGCCCCCTTCAAACCTTA CTTTGGAGGAAGTTACAGTGGGCCCAGTCTTTTGAAAATTTAATGGCTACAAAAATATGGTCCAGTAGT CTATGCAGCATATTTAAAAATGGAGCACAGTGACGAGAATATTCAATTCTGGATGGCATGTGAAACCTAT AAGAAAATTGCCTCACGGTGGAGCAGAATTTCTAGGGCAAAGAAGCTTTATAAGATTTACATCCAGCCAC AGTCCCCTAGAGAGATTAACATTGACAGTTCGACAAGAGAGACTATCATCAGGAACATTCAGGAACCCAC TGAAACATGTTTTGAAGAAGCTCAGAAAATAGTCTATATGCATATGGAAAGGGATTCCTACCCCAGATTT

CTAAAGTCAGAAATGTACCAAAAACTTTTGAAAAACTATGCAGTCCAACAACAGTTTC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC203804 protein sequence

Red=Cloning site Green=Tags(s)

MSRRNCWICKMCRDESKRPPSNLTLEEVLQWAQSFENLMATKYGPVVYAAYLKMEHSDENIQFWMACETY KKIASRWSRISRAKKLYKIYIQPQSPREINIDSSTRETIIRNIQEPTETCFEEAQKIVYMHMERDSYPRF

LKSEMYQKLLKTMQSNNSF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6310">https://cdn.origene.com/chromatograms/mk6310</a> g10.zip

**Restriction Sites:** Sgfl-Mlul



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_144766

ORF Size: 477 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 144766.3</u>

RefSeq Size:1538 bpRefSeq ORF:480 bpLocus ID:6003

UniProt ID: 014921



Cytogenetics: 1q31.2

**Protein Families:** Druggable Genome

**MW:** 19.1 kDa

**Gene Summary:** The protein encoded by this gene is a member of the regulator of G protein signaling (RGS)


family. RGS family members share similarity with S. cerevisiae SST2 and C. elegans egl-10 proteins, which contain a characteristic conserved RGS domain. RGS proteins accelerate GTPase activity of G protein alpha-subunits, thereby driving G protein into their inactive GDP-

bound form, thus negatively regulating G protein signaling. RGS proteins have been


implicated in the fine tuning of a variety of cellular events in response to G protein-coupled receptor activation. The biological function of this gene, however, is unknown. Two transcript

variants encoding the same isoform exist. [provided by RefSeq, Jul 2008]

## **Product images:**





Circular map for RC203804



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY RGS13 (Cat# RC203804, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RGS13 (Cat# [TA809778])(1:2000). Positive lysates [LY408107] (100ug) and [LC408107] (20ug) can be purchased separately from OriGene.







Western blot validation of overexpression lysate (Cat# [LY419008]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC208904] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified RGS13 protein (Cat# [TP303804]). The protein was produced from HEK293T cells transfected with RGS13 cDNA clone (Cat# RC203804) using MegaTran 2.0 (Cat# [TT210002]).