

## Product datasheet for RC203809L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Methionine Aminopeptidase 2 (METAP2) (NM\_006838) Human Tagged ORF Clone Lentiviral Particle

### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** Methionine Aminopeptidase 2 (METAP2) (NM\_006838) Human Tagged ORF Clone Lentiviral

Particle

**Symbol:** Methionine Aminopeptidase 2

**Synonyms:** MAP2; MNPEP; p67eIF2

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 006838

ORF Size: 1434 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC203809).

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 006838.3</u>

 RefSeq Size:
 3506 bp

 RefSeq ORF:
 1437 bp

 Locus ID:
 10988

 UniProt ID:
 P50579

 Cytogenetics:
 12q22

**Domains:** Peptidase\_M24





**Protein Families:** Druggable Genome, Protease


**MW:** 52.7 kDa

**Gene Summary:** The protein encoded by this gene is a member of the methionyl aminopeptidase family. The

encoded protein functions both by protecting the alpha subunit of eukaryotic initiation factor 2 from inhibitory phosphorylation and by removing the amino-terminal methionine residue from nascent proteins. Increased expression of this gene is associated with various forms of cancer, and the anti-cancer drugs fumagillin and ovalicin inhibit the protein by irreversibly binding to its active site. Inhibitors of this gene have also been shown to be effective for the treatment of obesity. A pseudogene of this gene is located on chromosome 2. Several transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Nov 2015]

### **Product images:**



[RC203809L3] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC203809L3V particle to overexpress human METAP2-Myc-DDK fusion protein.