

Product datasheet for **RC203813L3V**

BMP7 (NM_001719) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	BMP7 (NM_001719) Human Tagged ORF Clone Lentiviral Particle
Symbol:	BMP7
Synonyms:	OP-1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001719
ORF Size:	1293 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC203813).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001719.1
RefSeq Size:	4049 bp
RefSeq ORF:	1296 bp
Locus ID:	655
UniProt ID:	P18075
Cytogenetics:	20q13.31
Domains:	TGFb_propeptide, TGF-beta



[View online »](#)

Protein Families:	Adult stem cells, Cancer stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS, Induced pluripotent stem cells, Secreted Protein, Stem cell relevant signaling - TGFb/BMP signaling pathway
Protein Pathways:	Cytokine-cytokine receptor interaction, Hedgehog signaling pathway, TGF-beta signaling pathway
MW:	49.3 kDa
Gene Summary:	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer, which plays a role in bone, kidney and brown adipose tissue development. Additionally, this protein induces ectopic bone formation and may promote fracture healing in human patients. [provided by RefSeq, Jul 2016]