

## Product datasheet for **RC203882L3V**

### PCGF6 (NM\_001011663) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | PCGF6 (NM_001011663) Human Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | PCGF6  |
| Synonyms:                 | MBLR; RNF134   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_001011663   |
| ORF Size:                 | 1056 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC203882).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_001011663.1</a> , <a href="#">NP_001011663.1</a>  |
| RefSeq Size:              | 2248 bp  |
| RefSeq ORF:               | 1053 bp  |
| Locus ID:                 | 84108  |
| UniProt ID:               | <a href="#">Q9BYE7</a>   |
| Cytogenetics:             | 10q24.33   |
| Protein Families:         | Transcription Factors  |
| MW:                       | 39.2 kDa   |



[View online »](#)

**Gene Summary:**

The protein encoded by this gene contains a RING finger motif, which is most closely related to those of polycomb group (PcG) proteins RNF110/MEL-18 and BMI1. PcG proteins are known to form protein complexes and function as transcription repressors. This protein has been shown to interact with some PcG proteins and act as a transcription repressor. The activity of this protein is found to be regulated by cell cycle dependent phosphorylation. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]