

Product datasheet for RC203907L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Adiponectin Receptor 1 (ADIPOR1) (NM 015999) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Adiponectin Receptor 1 (ADIPOR1) (NM_015999) Human Tagged ORF Clone Lentiviral Particle

Symbol: Adiponectin Receptor 1

Synonyms: ACDCR1; CGI-45; CGI45; PAQR1; TESBP1A

NM 015999

Mammalian Cell

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ORF Size: 1125 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC203907).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 015999.2</u>

 RefSeq Size:
 2184 bp

 RefSeq ORF:
 1128 bp

 Locus ID:
 51094

 UniProt ID:
 Q96A54

 Cytogenetics:
 1q32.1

Domains: UPF0073

Protein Families: Druggable Genome, Transmembrane





Adiponectin Receptor 1 (ADIPOR1) (NM_015999) Human Tagged ORF Clone Lentiviral Particle – RC203907L1V

Protein Pathways: Adipocytokine signaling pathway

MW: 42.6 kDa

Gene Summary: This gene encodes a protein which acts as a receptor for adiponectin, a hormone secreted by

adipocytes which regulates fatty acid catabolism and glucose levels. Binding of adiponectin to the encoded protein results in activation of an AMP-activated kinase signaling pathway which affects levels of fatty acid oxidation and insulin sensitivity. A pseudogene of this gene is located on chromosome 14. Multiple alternatively spliced transcript variants have been found

for this gene. [provided by RefSeq, Mar 2014]