

## Product datasheet for RC203945L1

### beta IV Tubulin (TUBB4A) (NM\_006087) Human Tagged Lenti ORF Clone

#### Product data:

|                           |   |
|---------------------------|---|
| Product Type:             | Expression Plasmids   |
| Product Name:             | beta IV Tubulin (TUBB4A) (NM_006087) Human Tagged Lenti ORF Clone |
| Tag:                      | Myc-DDK   |
| Symbol:                   | beta IV Tubulin   |
| Synonyms:                 | beta-5; DYT4; TUBB4   |
| Mammalian Cell Selection: | None  |
| Vector:                   | pLenti-C-Myc-DDK (PS100064)                                       |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC203945).    |
| Restriction Sites:        | SgfI-MluI   |
| Cloning Scheme:           |   |

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


|           |           |
|-----------|-----------|
| ACCN:     | NM_006087 |
| ORF Size: | 1332 bp   |




[View online >](#)

|                               |   |
|-------------------------------|---|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>  |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol> |
| <b>RefSeq:</b>                | <a href="#">NM_006087.2</a>   |
| <b>RefSeq Size:</b>           | 2583 bp   |
| <b>RefSeq ORF:</b>            | 1335 bp   |
| <b>Locus ID:</b>              | 10382   |
| <b>UniProt ID:</b>            | <a href="#">P04350</a>  |
| <b>Cytogenetics:</b>          | 19p13.3   |
| <b>Domains:</b>               | tubulin   |
| <b>Protein Families:</b>      | Druggable Genome  |
| <b>Protein Pathways:</b>      | Gap junction, Pathogenic Escherichia coli infection   |
| <b>MW:</b>                    | 49.6 kDa  |
| <b>Gene Summary:</b>          | This gene encodes a member of the beta tubulin family. Beta tubulins are one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. Mutations in this gene cause hypomyelinating leukodystrophy-6 and autosomal dominant torsion dystonia-4. Alternate splicing results in multiple transcript variants encoding different isoforms. A pseudogene of this gene is found on chromosome X. [provided by RefSeq, Jan 2014]                                    |

Product images:



Circular map for RC203945L1



Double digestion of RC203945L1 using SgfI and MluI