

## Product datasheet for RC203969L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PLEKHO1 (NM\_016274) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: PLEKHO1 (NM\_016274) Human Tagged ORF Clone Lentiviral Particle

Symbol: PLEKHO1

**Synonyms:** CKIP-1; CKIP1; JBP; OC120

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_016274

ORF Size: 1227 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC203969).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 016274.3

 RefSeq Size:
 1627 bp

 RefSeq ORF:
 1230 bp

 Locus ID:
 51177

 UniProt ID:
 Q53GL0

 Cytogenetics:
 1q21.2

 Domains:
 PH

MW: 46.2 kDa







## **Gene Summary:**

Plays a role in the regulation of the actin cytoskeleton through its interactions with actin capping protein (CP). May function to target CK2 to the plasma membrane thereby serving as an adapter to facilitate the phosphorylation of CP by protein kinase 2 (CK2). Appears to target ATM to the plasma membrane. Appears to also inhibit tumor cell growth by inhibiting AKT-mediated cell-survival. Also implicated in PI3K-regulated muscle differentiation, the regulation of AP-1 activity (plasma membrane bound AP-1 regulator that translocates to the nucleus) and the promotion of apoptosis induced by tumor necrosis factor TNF. When bound to PKB, it inhibits it probably by decreasing PKB level of phosphorylation.[UniProtKB/Swiss-Prot Function]