

## Product datasheet for RC203986L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DP1 (TFDP1) (NM\_007111) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** DP1 (TFDP1) (NM\_007111) Human Tagged ORF Clone Lentiviral Particle

Symbol: DP1

Synonyms: DILC; Dp-1; DP1; DRTF1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_007111 **ORF Size:** 1230 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC203986).

OTI Disclaimer:

Sequence:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 007111.3

 RefSeq Size:
 2651 bp

 RefSeq ORF:
 1233 bp

 Locus ID:
 7027

 UniProt ID:
 Q14186

 Cytogenetics:
 13q34

**Protein Families:** Druggable Genome, Transcription Factors

E2F TDP





## DP1 (TFDP1) (NM\_007111) Human Tagged ORF Clone Lentiviral Particle - RC203986L2V

**Protein Pathways:** Cell cycle, TGF-beta signaling pathway

MW: 45.1 kDa

Gene Summary: This gene encodes a member of a family of transcription factors that heterodimerize with E2F

proteins to enhance their DNA-binding activity and promote transcription from E2F target genes. The encoded protein functions as part of this complex to control the transcriptional activity of numerous genes involved in cell cycle progression from G1 to S phase. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are found on

chromosomes 1, 15, and X.[provided by RefSeq, Jan 2009]