

Product datasheet for RC204041L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ERp72 (PDIA4) (NM_004911) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ERp72 (PDIA4) (NM_004911) Human Tagged ORF Clone Lentiviral Particle

Symbol: ERp72

Synonyms: ERp-72; ERP70; ERP72

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_004911

 ORF Size:
 1935 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC204041).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 004911.3</u>

 RefSeq Size:
 2970 bp

 RefSeq ORF:
 1938 bp

 Locus ID:
 9601

 UniProt ID:
 P13667

 Cytogenetics:
 7q36.1

Domains: thiored

Protein Families: Druggable Genome, Transmembrane





ERp72 (PDIA4) (NM_004911) Human Tagged ORF Clone Lentiviral Particle - RC204041L3V

Protein Pathways: Vibrio cholerae infection

MW: 72.9 kDa

Gene Summary: This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic

reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions.

The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence.

This protein, when bound to cyclophilin B, enhances the rate of immunoglobulin G intermolecular disulfide bonding and antibody assembly. [provided by RefSeq, Dec 2016]