

## Product datasheet for RC204056L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD99 (NM\_002414) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: CD99 (NM 002414) Human Tagged ORF Clone Lentiviral Particle

Symbol: CD99

Synonyms: HBA71; MIC2; MIC2X; MIC2Y; MSK5X

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 002414

ORF Size: 555 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC204056).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002414.3

RefSeq Size: 1255 bp

RefSeq ORF: 558 bp

Locus ID: 4267

UniProt ID: P14209

Cytogenetics: X;Y

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs), Leukocyte transendothelial migration





ORIGENE

**MW:** 18.8 kDa

**Gene Summary:** The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte

migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. There is a related pseudogene located immediately

adjacent to this locus. [provided by RefSeq, Mar 2016]