

Product datasheet for RC204093L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Retinoid X Receptor gamma (RXRG) (NM_006917) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Retinoid X Receptor gamma (RXRG) (NM_006917) Human Tagged ORF Clone Lentiviral Particle

Symbol: Retinoid X Receptor gamma

Synonyms: NR2B3; RXRC

Mammalian Cell None

Selection:

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_006917

ORF Size: 1389 bp

ORF Nucleotide

tide The ORF insert of this clone is exactly the same as(RC204093).

OTI Disclaimer:

Sequence:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006917.3</u>

 RefSeq Size:
 2205 bp

 RefSeq ORF:
 1392 bp

 Locus ID:
 6258

 UniProt ID:
 P48443

 Cytogenetics:
 1q23.3

HOLI, zf-C4





Retinoid X Receptor gamma (RXRG) (NM_006917) Human Tagged ORF Clone Lentiviral Particle – RC204093L1V

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Adipocytokine signaling pathway, Non-small cell lung cancer, Pathways in cancer, PPAR

signaling pathway, Small cell lung cancer, Thyroid cancer

MW: 50.9 kDa

Gene Summary: This gene encodes a member of the retinoid X receptor (RXR) family of nuclear receptors

which are involved in mediating the antiproliferative effects of retinoic acid (RA). This receptor forms dimers with the retinoic acid, thyroid hormone, and vitamin D receptors, increasing both DNA binding and transcriptional function on their respective response elements. This gene is expressed at significantly lower levels in non-small cell lung cancer cells. Alternatively

spliced transcript variants have been described. [provided by RefSeq, Jun 2010]